

# 计算机辅助几何设计


## 2021秋学期

Bézier Curves (continue)

陈仁杰

中国科学技术大学


# Recap



de Casteljau algorithm

Bernstein form

# Recap



## Bernstein form

Curve      basis function      control points

$$f(t) = \sum_{i=1}^n B_i(t)p_i$$

Useful properties for basis functions

- Smoothness
- Local control / support
- Affine invariance
- Convex hull property

# Degree elevation

- Given:  $b_0, \dots, b_n \rightarrow x(t)$
- Wanted:  $\bar{b}_0, \dots, \bar{b}_n, \bar{b}_{n+1} \rightarrow \bar{x}(t)$  with  $x = \bar{x}$
- Solution:

# Degree elevation

- Given:  $\mathbf{b}_0, \dots, \mathbf{b}_n \rightarrow x(t)$
- Wanted:  $\bar{\mathbf{b}}_0, \dots, \bar{\mathbf{b}}_n, \bar{\mathbf{b}}_{n+1} \rightarrow \bar{x}(t)$  with  $x = \bar{x}$
- Solution:
$$\begin{aligned}\bar{\mathbf{b}}_0 &= \mathbf{b}_0 \\ \bar{\mathbf{b}}_{n+1} &= \mathbf{b}_n \\ \bar{\mathbf{b}}_j &= \frac{j}{n+1} \mathbf{b}_{j-1} + \left(1 - \frac{j}{n+1}\right) \mathbf{b}_j \quad \text{for } j = 1, \dots, n\end{aligned}$$

# Proof

- Let's consider

$$\begin{aligned}(1-t)B_i^n(t) &= (1-t)\binom{n}{i}(1-t)^{n-i}t^i = \binom{n}{i}(1-t)^{n+1-i}t^i \\ &= \frac{n+1-i}{n+1}\binom{n+1}{i}(1-t)^{n+1-i}t^i \\ &= \frac{n+1-i}{n+1}B_i^{n+1}(t)\end{aligned}$$

Similarly

$$tB_i^n(t) = \frac{i+1}{n+1}B_i^{n+1}(t)$$

# Proof


Using results from

last slide

$$\begin{aligned}
 f(t) &= [(1-t) + t]f(t) = [(1-t) + t] \sum_{i=0}^n B_i^n(t) \mathbf{P}_i = \sum_{i=0}^n [(1-t)B_i^n(t) + tB_i^n(t)] \mathbf{P}_i \\
 &= \sum_{i=0}^n \left[ \frac{n+1-i}{n+1} B_i^{n+1}(t) + \frac{i+1}{n+1} B_{i+1}^{n+1}(t) \right] \mathbf{P}_i = \sum_{i=0}^n \frac{n+1-i}{n+1} B_i^{n+1}(t) \mathbf{P}_i + \sum_{i=0}^n \frac{i+1}{n+1} B_{i+1}^{n+1}(t) \mathbf{P}_i \\
 &= \sum_{i=0}^n \frac{n+1-i}{n+1} B_i^{n+1}(t) \mathbf{P}_i + \sum_{i=1}^{n+1} \frac{i}{n+1} B_i^{n+1}(t) \mathbf{P}_{i-1} \\
 &= \sum_{i=0}^{n+1} \frac{n+1-i}{n+1} B_i^{n+1}(t) \mathbf{P}_i + \sum_{i=0}^{n+1} \frac{i}{n+1} B_i^{n+1}(t) \mathbf{P}_{i-1} \\
 &= \sum_{i=0}^{n+1} B_i^{n+1}(t) \left[ \frac{n+1-i}{n+1} \mathbf{P}_i + \frac{i}{n+1} \mathbf{P}_{i-1} \right]
 \end{aligned}$$


Adding null terms,  $i = n+1, i = 0$

# Degree elevation: Example




- $\bar{b}_0 = b_0$        $\bar{b}_j = \frac{j}{n+1} b_{j-1} + \left(1 - \frac{j}{n+1}\right) b_j$
- $\bar{b}_{n+1} = b_n$        $j = 1, \dots, n$

# Degree elevation: Example




- $\bar{b}_0 = b_0$        $\bar{b}_j = \frac{j}{n+1} b_{j-1} + \left(1 - \frac{j}{n+1}\right) b_j$
- $\bar{b}_{n+1} = b_n$        $j = 1, \dots, n$

# Degree elevation: Example




- $\bar{b}_0 = b_0$       
$$\bar{b}_j = \frac{j}{n+1} b_{j-1} + \left(1 - \frac{j}{n+1}\right) b_j$$
- $\bar{b}_{n+1} = b_n$       
$$j = 1, \dots, n$$

# Degree elevation: Example



- $\bar{b}_0 = b_0$        $\bar{b}_j = \frac{j}{n+1} b_{j-1} + \left(1 - \frac{j}{n+1}\right) b_j$
- $\bar{b}_{n+1} = b_n$        $j = 1, \dots, n$


# Degree elevation: Example



- $\bar{b}_0 = b_0$
- $\bar{b}_{n+1} = b_n$
- $$\bar{b}_j = \frac{j}{n+1} b_{j-1} + \left(1 - \frac{j}{n+1}\right) b_j$$
$$j = 1, \dots, n$$


$$\frac{1}{4}$$
$$\frac{3}{4}$$

# Degree elevation: Example




- $\bar{b}_0 = b_0$        $\bar{b}_j = \frac{j}{n+1} b_{j-1} + \left(1 - \frac{j}{n+1}\right) b_j$
- $\bar{b}_{n+1} = b_n$        $j = 1, \dots, n$

# Degree elevation: Example




- $\bar{b}_0 = b_0$        $\bar{b}_j = \frac{j}{n+1} b_{j-1} + \left(1 - \frac{j}{n+1}\right) b_j$
- $\bar{b}_{n+1} = b_n$        $j = 1, \dots, n$

# Degree elevation



For repeated degree elevation, the Bézier polygon converges to the Bézier curve. (slow convergence)

# Degree elevation



- $\bar{b}_0 = b_0$        $\bar{b}_j = \frac{j}{n+1} b_{j-1} + \left(1 - \frac{j}{n+1}\right) b_j$
- $\bar{b}_{n+1} = b_n$        $j = 1, \dots, n$

# Bézier Curves

## Subdivision

# Subdivision


- **Given:**  $b_0, \dots, b_n \rightarrow x(t), t \in [0,1]$

- **Wanted:**  $b_0^{(1)}, \dots, b_n^{(1)} \rightarrow x^{(1)}(t),$

- $b_0^{(2)}, \dots, b_n^{(2)} \rightarrow x^{(2)}(t),$


**with**  $x = x^{(1)} \cup x^{(2)}$

# Subdivision: Example



de Casteljau scheme

# Subdivision: Example




de Casteljau scheme


# Subdivision

Solution:  $b_i^{(1)} = b_0^i, b_i^{(2)} = b_0^{n-i}$  for  $i = 0, \dots, n$

That means that the new points are intermediate points of the de Casteljau algorithm!




# Curve range




parameterization:  $t \in [0, 1]$

# Curve range



# Summary & Outlook

- **Bézier curves and curve design**
  - The rough form is specified by the position of the control points
  - Results: smooth curve approximating the control points
  - Computation / Representation:
    - de Casteljau algorithm
    - Bernstein form
- Problems:
  - High polynomial degree
  - Moving a control point can change the whole curve
  - Interpolation of points
  - →**Bézier splines**



# **Matrix representations**

**(common in software implementations)**

# Homogeneous coordinates

$$[P] = \begin{pmatrix} x_1 & y_1 & z_1 & 1 \\ x_2 & y_2 & z_2 & 1 \\ x_3 & y_3 & z_3 & 1 \\ \dots & \dots & \dots & \dots \\ x_n & y_n & z_n & 1 \end{pmatrix}$$

# Transformations

- Basic representation  $[P^*] = [P][T]$ 
  - $[P^*]$  is the new coordinates matrix
  - $[P]$  is the original coordinates matrix, or points matrix
  - $[T]$  is the transformation matrix

# Transformations

- Translation (2D example)

$$[T_t] = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ x & y & 0 & 1 \end{pmatrix}$$

$$[P^*] = [P][T_t]$$

# Transformations

- Basic representation  $[P^*] = [P][T]$ 
  - $[P^*]$  is the new coordinates matrix
  - $[P]$  is the original coordinates matrix, or points matrix
  - $[T]$  is the transformation matrix

$$[P] = \begin{pmatrix} x_1 & y_1 & 0 \\ x_2 & y_2 & 0 \\ x_3 & y_3 & 0 \\ \dots & \dots & \dots \\ x_n & y_n & 0 \end{pmatrix}$$

# Transformations

- Uniform scaling

$$[T] = \begin{pmatrix} s & 0 & 0 & 0 \\ 0 & s & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

- Non-uniform scaling

$$[T] = \begin{pmatrix} s_x & 0 & 0 & 0 \\ 0 & s_y & 0 & 0 \\ 0 & 0 & s_z & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

# Transformations

- Rotation 2D

$$\left. \begin{array}{l} x = r \cos \alpha \\ y = r \sin \alpha \end{array} \right\} \text{Original coordinates of point } P$$

$$\left. \begin{array}{l} x^* = r \cos(\alpha + \theta) \\ y^* = r \sin(\alpha + \theta) \end{array} \right\} \text{The new coordinates}$$

$$[x^* \quad y^* \quad 0 \quad 1] = [x \quad y \quad 0 \quad 1] \begin{pmatrix} \cos \theta & \sin \theta & 0 & 0 \\ -\sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

# Transformations

- Rotation about an arbitrary axis (2D)
  - Translate the fixed axis so it coincides with z-axis
    - apply to object
  - Rotate object about the axis
  - Translate object back

$$[P^*] = [P][T_t][T_r][T_{-t}]$$

# Transformations

- Rotation about an arbitrary axis (2D)

Step 1: Translate the fixed axis so it coincides with z-axis

Step 2: Rotate object about the axis

Step 3: Translate the fixed axis back to the original position

$$[P^*] = [P][T_t][T_r][T_{-t}]$$

# Transformations

- Scaling with an arbitrary point  $(x, y)$

$$[P^*] = [P][T_t][T_s][T_{-t}]$$

$$= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -x & -y & 0 & 1 \end{pmatrix} \begin{pmatrix} s & 0 & 0 & 0 \\ 0 & s & 0 & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ x & y & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} s & 0 & 0 & 0 \\ 0 & s & 0 & 0 \\ 0 & 0 & s & 0 \\ x - sx & y - sy & 0 & 1 \end{pmatrix}$$

# Transformations

- Rotation about an arbitrary point  $(x, y)$


$$[T_{\text{cond}}] = [T_t][T_s][T_{-t}]$$

$$= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -x & -y & 0 & 1 \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta & 0 & 0 \\ -\sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ x & y & 0 & 1 \end{pmatrix}$$

# Transformations

- Mirroring about x-axis (negative scaling along y-axis)

$$[P^*] = [2 \quad 2 \quad 0 \quad 1] \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$
$$= [2 \quad -2 \quad 0 \quad 1]$$



# Transformations

- Mirroring about arbitrary axis
  - Translate line to pass through origin
  - Rotate axis to coincide with  $x$ -axis
  - Mirror about  $x$ -axis
  - Rotate back
  - Translate back to original position

$$[P^*] = [P][T_t][T_r][T_m][T_{-r}][T_{-t}]$$

# Transformations

- Rotation about coordinates axes (3D)

$$[T_{rz}] = \begin{pmatrix} \cos \theta & \sin \theta & 0 & 0 \\ -\sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$[T_{rx}] = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$[T_{ry}] = \begin{pmatrix} \cos \theta & 0 & -\sin \theta & 0 \\ 0 & 1 & 0 & 0 \\ \sin \theta & 0 & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

# Transformations

- Rotation  $\theta$  about an arbitrary axis (3D)
  1. Translate the given line so that it will pass through the origin
  2. Rotate about the  $x$ -axis so that the line lies in the  $xz$ -plane (angle  $\alpha$ )
  3. Rotate about the  $y$ -axis so that the line coincides with the  $z$ -axis (angle  $\phi$ )
  4. Rotate the geometric object about the  $z$ -axis (angle  $\theta$  – given rotation angle)
  5. Reverse of step 3
  6. Reverse of step 2
  7. Reverse of step 1

$$[P^*] = [P][T_t][T_r]_\alpha[T_r]_\phi[T_r]_\theta[T_r]_{-\phi}[T_r]_{-\alpha}[T_{-t}]$$

Alternatively you can use Quaternions!

# Bézier Curves

- Cubic Bézier curves

$$f(t) = P_0 B_0^{(3)} + P_1 B_1^{(3)} + P_2 B_2^{(3)} + P_3 B_3^{(3)}$$

$$B_0^{(3)}(t) = \frac{3!}{0! 3!} t^0 (1-t)^3 = (1-t)^3$$

$$B_1^{(3)} = \frac{3!}{1! 2!} t^1 (1-t)^2 = 3t(1-t)^2$$

$$B_2^{(3)} = \frac{3!}{2! 1!} t^2 (1-t)^1 = 3t^2(1-t)$$

$$B_3^{(3)} = \frac{3!}{3! 0!} t^3 (1-t)^0 = t^3$$

# Bézier Curves

- Cubic Bézier curves

$$f(t) = P_0 B_0^{(3)} + P_1 B_1^{(3)} + P_2 B_2^{(3)} + P_3 B_3^{(3)}$$

- In Matrix form:

- -The curve

$$f(t) = [t^3 \quad t^2 \quad t \quad 1] \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{bmatrix} P_0 \\ P_1 \\ P_2 \\ P_3 \end{bmatrix}$$

- -The tangent

$$f'(t) = [3t^2 \quad 2t \quad 1 \quad 0] \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{bmatrix} P_0 \\ P_1 \\ P_2 \\ P_3 \end{bmatrix}$$

# B-spline Curves (to be covered later)

- Uniform cubic B-Spline curve

$$f_i(t) = \frac{1}{6} [t^3 \quad t^2 \quad t \quad 1] \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 0 & 3 & 0 \\ 1 & 4 & 1 & 0 \end{pmatrix} \begin{bmatrix} P_{i-1} \\ P_i \\ P_{i+1} \\ P_{i+2} \end{bmatrix}$$

# B-spline Curves (to be covered later)

- Other splines:

- Catmull-Rom

$$f_i(t) = [t^3 \quad t^2 \quad t \quad 1] \frac{1}{2} \begin{pmatrix} -1 & 3 & -3 & 1 \\ 2 & -5 & 4 & -1 \\ -1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \begin{bmatrix} P_{i-1} \\ P_i \\ P_{i+1} \\ P_{i+2} \end{bmatrix}$$

- Cardinal splines

$$\begin{pmatrix} -a & 2-a & a-2 & a \\ 2a & a-3 & 3-2a & -a \\ -a & 0 & a & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

- Tensioned splines

$$\frac{1}{6} \begin{pmatrix} -a & 12-9a & 9a-12 & a \\ 2a & a-3 & 18-15a & -a \\ -3a & 0 & 3a & 0 \\ 0 & 6-2a & a & 0 \end{pmatrix}$$